

Calf Pneumonia Vaccines & Regimes

Vaccine	What Covered	Administration	Age 1 st Dose	Repeat Dose
Bovilis RSP	RSV, PI3, Pasteurella	subcutaneous	>2weeks	4weeks later
Bovilis IBR (live)	IBR	intranasal (preferably)	>4weeks single dose	Annual booster
		intramuscular	>4weeks single dose	Annual booster
Bovilis IBR marker (live)	IBR	intranasal	>2 weeks	Repeat at 3-4 months if first dose given before 3 months Booster every 6 months
		intramuscular	>3 months	
Imuresp	IBR, PI3	intranasal	>3weeks	Calves vaccinated before 10 weeks require second dose after 10 weeks. Booster every 6 months
Pastobov	Pasteurella	i/m or s/c injection	>4weeks	3-4 weeks later. Annual booster
Rispoval 3	PI3, RSV, BVD,	intramuscular	>3months	3-4 weeks later. Booster 6 months
Rispoval 4	PI3, RSV, BVD, IBR	intramuscular	>3weeks	3-4 weeks later. If first dose given before 3 months a third dose is given after 3 months.
Rispoval IBR Marker (atten)	IBR	subcutaneous	>3months	3-5 weeks later. Calves injected under 3 months need revaccinating after 3 months. Booster in 3 months.
Rispoval IBR Marker (live)	IBR	intranasal	>2weeks	3-5 weeks later intramuscular
		intramuscular	>3months	3-5 weeks later if seronegative
Rispoval Pasteurella	pasteurella	intramuscular	>3 months single dose	
Rispoval RS	RSV	intramuscular	>7days	3-4 weeks later. Calves injected under 4 months require third injection after 4months of age
Rispoval RS &PI3 I/N	PI3, RSV	intranasal	>3weeks single dose	
Tracherine	IBR	intranasal	>3weeks	If vaccinated under 10 weeks calves will require another dose after 10 weeks